

Phase structure of $(2 + 1)d$ strongly coupled lattice gauge theories

Costas G. Strouthos^a

^aDepartment of Physics, Box 90305, Duke University, Durham, North Carolina 27708, USA.

We study the chiral phase transition in $(2 + 1)d$ strongly coupled $U(N)$ lattice gauge theories with staggered fermions. We show with high precision simulations performed directly in the chiral limit that these models undergo a Berezinski-Kosterlitz-Thouless (BKT) transition. We also show that this universality class is unaffected even in the large N limit.

1. Introduction

The behavior of symmetries at finite temperature is one of the most outstanding problems in field theory. In recent years we have witnessed revised interest in the chiral phase transition in QCD. The problem of symmetry breaking and its restoration is intrinsically non-perturbative and therefore most of our knowledge about the phenomenon comes from lattice simulations. However, computing quantities in lattice QCD with massless quarks is a notoriously difficult problem, because most known algorithms break down in the chiral limit. In addition, the simulations near T_c must be performed on lattices with large spatial sizes in order to control large finite size effects due to the diverging correlation length.

A useful simplification of QCD occurs in the strong coupling limit, which retains much of the underlying physics of QCD except for large lattice artifacts. In this limit chiral symmetry breaking and its restoration at finite temperature have been studied using large N and large d expansions [1]. However, since these approaches are based on mean field analysis they cannot help in determining the universality of phase transitions.

Interestingly, lattice QCD with one staggered fermion interacting with $U(N)$ gauge fields can be mapped into a monomer-dimer system in the strong coupling limit [2]. Here, we present numerical results for the finite temperature critical behavior of the $(2 + 1)d$ model. Our data were generated with a recently developed very efficient cluster algorithm [3], which allows us to

perform precision calculations in the chiral limit. In agreement with expectations from universality and dimensional reduction, we show convincingly [4] that the chiral phase transition belongs to the BKT universality class [5].

The partition function of the model we study here is given by

$$Z = \sum_{[n,b]} \prod_{\langle ij \rangle} (z_{\langle ij \rangle})^{b_{\langle ij \rangle}} \frac{(N - b_{\langle ij \rangle})!}{b_{\langle ij \rangle}! N!} \prod_i \frac{N!}{n_i!} m^{n_i}, \quad (1)$$

and is discussed in detail in [2,3]. Here $n_i = 0, 1, 2, \dots, N$ refers to the number of monomers on the site i , $b_{\langle ij \rangle} = 0, 1, 2, \dots, N$ represents the number of dimers on the bond $\langle ij \rangle$, m is the monomer weight, $z_{\langle ij \rangle} = \eta_{ij}^2$ are the dimer weights. Note that while spatial dimers carry a weight 1, temporal dimers carry a weight T . The sum is over all monomer-dimer configurations $[n, b]$ which are constrained such that the sum of the number of monomers at each site and the dimers that touch the site is always N (the number of colors). In this work we choose $L_x = L_y = L$. One can study the thermodynamics of the model by working on asymmetric lattices with $L_t \ll L$ and allowing T to vary continuously.

2. Results

In this section we present the results for the $T \neq 0$ critical behavior of the model. The observables used in this work are the chiral susceptibility χ_c and the winding number susceptibility χ_w . The latter is proportional to the helicity modulus and describes the response of the system to

Table 1

Fits for χ_c and χ_w near T_c .

T	η	$2r$	$\chi_1^2/\text{d.o.f}$	$1/(2c)$	$\chi_2^2/\text{d.o.f}$
1.00	0.222(5)	0	0.2	0.2343(8)	183.2
1.02	0.235(5)	-0.3(2)	1.5	0.2411(5)	53.4
1.04	0.251(5)	-0.07(2)	0.4	0.2483(5)	2.8
1.06	0.249(5)	-0.03(2)	0.5	0.2583(5)	33.8
1.10	0.388(5)	-0.12(2)	3.6	0.2831(5)	770.0
1.14	0.569(6)	-1.24(3)	480	—	—

a perturbation that distorts the direction of the spontaneous magnetization. The winding number susceptibility has been used successfully in to demonstrate BKT behavior in other models [6]. We fixed $L_t = 4$ and computed χ_c and χ_w as a function of L . If T_c is the critical temperature, then the BKT theory predicts


$$\chi_c \propto \begin{cases} L^{2-\eta(T)} & T < T_c \\ L^{1.75} [\log(L)]^{0.125} & T = T_c \\ \text{constant} & T > T_c \end{cases} \quad (2)$$

and


$$\chi_w = \begin{cases} 1/[2\eta(T)] + \alpha_1 L^{-\alpha_2} & T < T_c \\ [2 + 1/\log(L/L_0)] & T = T_c \\ \alpha_3 \exp(-\alpha_4 L) & T > T_c \end{cases} \quad (3)$$

in the large L limit. The critical exponent $\eta(T)$ is expected to change continuously with T , but remains in the range $0 \leq \eta(T) < 0.25$. In order to confirm these predictions we computed χ_c and χ_w for lattices ranging from $L = 32$ to $L = 750$ and for $N = 1, \dots, 32$.

Let us first discuss our results for $N = 1$. In figures 1 and 2 we plot χ_c and χ_w as functions of L for different values of T . We find that χ_c fits well to the form $bL^{2-\eta}(\log(L))^{-2r}$ when $T \leq 1.06$. We also find that the logarithmic term is unimportant for $T \leq 1.0$, whereas at $T = 1.04$ the value $2r = 0.7(2)$ is close to the BKT prediction which is 0.125. The values of η , $2r$ and the quality of the fits $\chi_1^2/\text{d.o.f}$ are shown in table 1. We also fit the data for χ_w to the form $(c + 1/\log(L/L_0))$. The values of $1/(2c)$ are also shown in the table. Finally, using the fact that $\chi_w = (2 + 1/\log(L/L_0))$ is exactly valid at T_c we fit the data for χ_w to this form for various values of T . The values of $\chi_2^2/\text{d.o.f}$ are shown in the last column of table

Figure 1. Plot of χ_c vs. L for various T .

1. Based on where the minimum in $\chi_2^2/\text{d.o.f}$ occurs we estimate $T_c = 1.040(5)$. Our value of $\eta = 0.250(5)$ at T_c is in excellent agreement with the BKT prediction.

Figure 2. Plot of χ_w vs. L for various T .




Figure 3. Plot of η as a function of N for $T = 1.0$. The data fit very well to the form $\eta = 0.169(6)/N + 0.050(9)/N^2$ with $\chi^2/\text{d.o.f} = 1.2$.

We also checked that χ_c and χ_w show similar evidence for a BKT transition at larger values of N . Using techniques similar to the ones we used for $N = 1$ we computed T_c for various values of N . We find that $T_c = 0.708(6)N + 1.40(4) - 1.07(4)/N$ fits our results very well for all values of N with a $\chi^2/\text{d.o.f}$ of 1.1. The dependence of the coefficients of this polynomial on L_t is still under investigation.




Figure 4. Plot of η as a function of T/T_c for various values of N .

With regards to the N dependence of our results we find two interesting observations. Witten argued that when the symmetry is $U(1)$ the large N analysis is still applicable since $\eta \sim 1/N$ at large N [7]. Our results for a fixed $T = 1.0$, shown

in figure 3, do agree with his conjecture. Interestingly, as N becomes large and T/T_c is held fixed instead of T , we find that $\eta \neq 0$ even in the large N limit. Figure 4 shows that η approaches an interesting function of T/T_c as N becomes large. Extending this observation to QCD, we think that the t'Hooft limit (large N with g^2N held fixed) may be quite similar [8].

3. Summary

We presented high precision results from simulations of $(2+1)d$ $U(N)$ strongly coupled lattice gauge theories at $T \neq 0$ and we showed convincing evidence that the models undergo a BKT phase transition. In addition, we showed that this universality class is unaffected even in the large N limit, implying that the mean field analysis often used in this limit breaks down in the critical region.

Acknowledgements

This work was done in collaboration with Shailesh Chandrasekharan and it was supported in part by the NSF grant DMR-0103003 and DOE grant DE-FG-96ER40945.

REFERENCES

1. N. Kawamoto and J. Smit, Nucl. Phys. **B192** (1981) 100; H. Klumberg-Stern, A. Morel and B. Petersson, Nucl. Phys. **B215** (1983) 527; P.H. Damgaard, N. Kawamoto and K. Shigemoto, Phys. Rev. Lett. **53** (1984) 527.
2. P. Rossi and U. Wolff, Nucl. Phys. **B248** (1984) 105.
3. D.H. Adams and S. Chandrasekharan, Nucl. Phys. **B662** (2003) 220.
4. S. Chandrasekharan and C.G. Strouthos, hep-lat/0306034.
5. V.L. Berezinski, Sov. Phys. JETP **34** (1971) 610; J.M. Kosterlitz and D.J. Thouless, J. Phys. **C6** (1973) 1181.
6. K. Harada and N. Kawashima, J. Phys. Soc. Jpn. **67** (1998) 2768; S. Chandrasekharan and J.C. Osborn, Phys. Rev. **B66** (2002) 045113.
7. E. Witten, Nucl. Phys. **B145** (1978) 110.
8. G. t'Hooft, Nucl. Phys. **B72** (1974) 461.